# Mindfulness-based treatments for co-occurring depression and substance use disorders: what can we learn from the brain?

## Judson A. Brewer<sup>1</sup>, Sarah Bowen<sup>2</sup>, Joseph T. Smith<sup>1</sup>, G. Alan Marlatt<sup>2</sup> & Marc N. Potenza<sup>1,3</sup>

Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA,<sup>1</sup> Department of Psychology, University of Washington, Seattle, WA, USA<sup>2</sup> and Child Study Center, Yale University School of Medicine, New Haven, CT, USA<sup>3</sup>

## ABSTRACT

Both depression and substance use disorders represent major global public health concerns and are often co-occurring. Although there are ongoing discoveries regarding the pathophysiology and treatment of each condition, common mechanisms and effective treatments for co-occurring depression and substance abuse remain elusive. Mindfulness training has been shown recently to benefit both depression and substance use disorders, suggesting that this approach may target common behavioral and neurobiological processes. However, it remains unclear whether these pathways constitute specific shared neurobiological mechanisms or more extensive components universal to the broader human experience of psychological distress or suffering. We offer a theoretical, clinical and neurobiological perspective of the overlaps between these disorders, highlight common neural pathways that play a role in depression and substance use disorders. Finally, we discuss how advances in our understanding of potential mechanisms of mindfulness training may offer not only unique effects on depression and substance use, but also offer promise for treatment of co-occurring disorders.

**Keywords** Addiction, co-occurring disorders, depression, dual diagnosis, functional magnetic resonance imaging (fMRI), mindfulness, substance use treatment.

Correspondence to: Judson Brewer, VA Connecticut Healthcare System, 950 Campbell Avenue, Building 36, Room 142, West Haven, CT 06516, USA. E-mail: judson.brewer@yale.edu

Submitted 22 May 2009; initial review completed 22 July 2009; final version accepted 16 November 2009

# MINDFULNESS TRAINING MAY PROVE EFFECTIVE FOR CO-OCCURRING DEPRESSION AND SUBSTANCE USE DISORDERS WHERE OTHER TREATMENTS FAIL

Mindfulness training (MT) may target common underlying mechanisms of major depressive disorder (MDD) and substance use disorders (SUDs), providing an effective treatment for co-occurrence of these maladies where few currently exist. We will use 'co-occurring disorder' and 'dual diagnosis' to refer to the co-occurrence of MDD and an SUD in which 'diagnoses of these disorders must occur simultaneously or within a one year time frame of each other' [1]. As first-line treatment for substance-induced mood disorders is to treat the underlying SUD [2], substance-induced mood disorders will not be included in this discussion. Additionally, although co-occurring psychotic and anxiety disorders are common and warrant discussion, the majority of the current discussion will be limited to co-occurring MDD and SUDs.

# MDD AND SUDS: SIGNIFICANT AND GROWING PROBLEMS

Both MDD and SUDs are major public health problems. Overall, unipolar depressive disorders were the fourth leading cause of disease burden in 2002, and are projected to be the second leading cause by 2030 [3]. Further, MDD and SUDs have been found to co-occur frequently: about one-third of individuals with MDD also have symptoms consistent with an SUD [4], and life-time prevalence of a co-occurring SUD ranges from 30% to 42.8% [3,5]. From 1992–2002, the rates of pastyear major depressive episodes among people with a concurrent SUD increased from 10% to 15% [6]. These data highlight the large and growing burden of comorbid MDD and SUDs.

There are currently multiple empirically supported behavioral treatments for both MDD and SUDs as individual disorders. Well-supported behavioral treatments for depression include cognitive behavioral therapy (CBT). behavioural-activation therapy, mindfulness-based cognitive therapy and exercise [7]. Similarly, substance abuse treatments include CBT, relapse prevention, motivational interviewing (MI), contingency management (CM) and 12-Step programs [8-10]. Despite significant research on treating these disorders individually, few treatments have been evaluated for individuals with co-occurring MDD and SUDs [11]. MI has demonstrated efficacy for many types of substance abuse, although few studies have been conducted with dual diagnosis patients [12,13]. CBT has been modified for people with SUDs and depression and has shown moderate effects therein [14]. CM has been adopted more recently for dually diagnosed populations, although mood symptomatology was not measured in these studies, leaving its broader efficacy unverified [13].

# MINDFULNESS TRAINING FOR CO-OCCURRING DISORDERS: THEORETICAL, CLINICAL AND NEUROBIOLOGICAL PERSPECTIVES

During the past two decades, the emergence of studies on treatments incorporating mindfulness training (MT) has offered promise for the treatment of MDD and SUDs independently. We offer a theoretical, behavioral and neurobiological exploration of MT for treatment of these disorders when they occur concurrently.

### Theoretical perspectives

A recent consensus definition of mindfulness emphasizes two complementary elements: (1) the placement of attention on the immediate experience; and (2) adopting an open, curious, accepting attitude towards that experience [15]. It is believed that although the capacity for mindfulness is inherent, the majority of individuals move through life on 'auto-pilot,' performing daily activities based on habitual behavioral patterns while their minds are elsewhere [16]. Mindfulness is developed through a continual practice of 'awakening' to present-moment experiences. Traditionally, this is taught through meditation practices that focus first on developing concentration capacities by bringing attention to an object repeatedly, such as the breath, then broadening the attention to include all physical and mental events that are experienced (e.g. bodily sensations, emotions and thoughts). Instruction for mindfulness practice can be as simple as:

'when sitting, know you are sitting; when thinking, know you are thinking' [J. Goldstein, personal communication]. The intention is to bring a non-judgemental, objective and accepting quality to this observation.

### **Clinical perspectives**

While mindfulness practices are centuries old, they have only recently become popular in western medicine and psychology. One of the earliest incorporations of mindfulness into medical treatment paradigms was through the mindfulness-based stress reduction (MBSR) program at the University of Massachusetts [17]. MBSR showed efficacy in the treatment of chronic pain where other treatments had failed [18]. After the initial success in treating chronic pain, the effectiveness of mindfulnessbased therapies was investigated for treatment of other conditions including anxiety disorders [19-22], addiction [23-26] and depression [27,28], although methodological quality of these studies has been suboptimal [29]. Mindfulness-based cognitive therapy (MBCT), based largely upon MBSR, has shown an absolute reduction of 44–50% in the relapse rate for individuals with three or more episodes of depression both in initial and replication studies [27,28]. A study of incarcerated, substanceabusing individuals who were taught mindfulness meditation revealed significant reduction in substance use 3 months following release from incarceration, as well as reductions in anxiety and depression [30]. Another study of individuals with alcohol and cocaine use disorders suggested that MT may be as effective as CBT in preventing relapse with specific effects on psychological and physiological stress pathways [24]. A further study of a mindfulness-based treatment for substance use disorders suggested that participation in treatment was associated with greater decreases in craving and substance use compared to a treatment-as-usual method [31], and that the treatment may lessen the relation between depressive symptomotology and craving, thereby decreasing substance use. Taken together, these data provide a rationale for the hypothesis that MT may target shared underlying mechanisms in MDD and SUDs in a dually diagnosed population.

#### Neurobiological perspectives

Evidence for common underlying neural pathways in MDD and SUDs may explain some of the shared mechanisms in these disorders, offering a useful perspective on potential targets of treatment for dually diagnosed individuals. Multiple approaches have investigated the pathophysiology of MDD and SUDs, including those based on genetics, neurotransmitters and endocrine systems (reviewed elsewhere [32–35]). Regional brain activation studies have provided unique insight into the pathophysi-

ology of both MDD and SUDs. Many regions have shown overlap between MDD and SUDs, suggesting possible mutual underlying pathophysiologies. For example, regions of the ventromedial prefrontal cortex (vmPFC), which is important for homeostasis, emotional regulation and decision-making, have shown dysfunction in both MDD [36-40] and SUDs [41-47]. The dorsolateral PFC (dlPFC), which is involved in working memory, attention, initiation of cognitive control and conflict-induced behavioral adjustment [48-50], the amygdala, which contributes to the formation and storage of memories associated with emotional events, memory consolidation and reward learning and motivation [51-53] and the insula, which contributes to 'sensing' of somatic states through its representation of bodily sensation [54], have all shown importance in both MDD and SUDs [36,39,55-62]. While these data suggest overlapping neuroanatomical correlates for MDD and SUDs that may be reflected in common intermediary phenotypes or endophenotypes, they do not demonstrate functional commonalities directly.

The psychological and behavioral correlates of brain function in MDD and SUDs may provide insight into potential targets of effective treatments. For example, several studies suggest that rumination and stress are seen commonly in both MDD and SUDs. Rumination, described as self-focused attention on symptoms of distress without engagement in active problem solving [63], is conceived of as an automatic behavior often acquired during a first depressive episode [27]. People who engage in rumination when distressed are more likely to become depressed and have longer periods of depression [63]. Similarly, temporary distraction, which may break the ruminative cycle, may lead to a decrease in dysphoria [64]. Brain regions that are implicated in rumination include the amygdala and PFC [63]. Ruminators have shown lower PFC activity than controls when attempting to inhibit negative distracters, as well as difficulty with cognitive shifting [65]. These data are suggestive of dysfunctional cognitive control circuitry [66]. Other studies suggest that rumination is associated with increased amygdalar activity during processing of emotional stimuli [67,68]. A role for rumination in SUDs has been suggested by work showing that a tendency to ruminate is associated with greater inclinations to use alcohol or other substances [69]. Although preliminary, these data suggest that dysfunction in specific brain regions may correlate with rumination and contribute to habitual behaviors in both MDD and SUDs.

Links between stress, depression and substance use have also been established (reviewed elsewhere [70,71]). For example, exposure to stressful life events has been associated repeatedly with MDD [32,71]. Also, depressed patients often exhibit elevated plasma cortisol (the dominant circulating stress glucocorticoid hormone in humans) and abnormal cortisol suppression [72]. Similarly, stress has been shown to be instrumental in SUDs: stress cross-sensitizes to both stimulant [73] and alcohol [74] use, induces craving [75] and increases selfadministration of drugs such as amphetamines [76], cocaine [77] and alcohol [78]. This is influenced probably by stress hormones, as in healthy volunteers stimulants induce cortisol release, while the magnitude of the reported subjective 'high' correlates with plasma cortisol concentrations [79].

The above data suggest that MDD and SUDs share several phenotypes such as stress vulnerability and rumination, pointing at possible mutual underlying neurobiological dysfunctions. Still unclear are the details of how specific this overlap may be. However, these commonalities may provide a sufficient framework from which to develop treatments targeted at shared brain and behavioral dysfunction.

## MECHANISM OF ACTION IN MINDFULNESS TRAINING

Recent neurobiological, cognitive and behavioral data support two specific components of mindfulness, attention and acceptance that may target the common intermediary phenotypes of rumination and stress directly, highlighting their potential utility in the treatment of MDD and SUDs (reviews of the general mechanisms of mindfulness can be found elsewhere [15,80–83]).

#### Attention

#### Theoretical perspectives

One primary aim of MT is to shift attention from a passive, wandering state ('default mode'—see below) to an active, intentional state. For example, during concentration meditation, when an individual's mind strays from the object of attention or is distracted by other stimuli, the individual is instructed to 'bring the attention back' intentionally to the intended focus. With practice, individuals retrain their minds to more continually 'pay attention, on purpose, in the present moment' [17].

#### Clinical perspectives

The link between attention training and treatment of depression and addiction may not be apparent initially. It has been hypothesized that focusing and sustaining attention on present experience increases the ability to notice overlearned behavioral patterns as they arise, allowing for individualized interventions (whether cognitive or behavioral) that interrupt these patterns [15,84]. For example, early in major depressive episodes, associa-

tions between mood and depressive thought patterns are established and can be reactivated in periods of dysphoria [85]. In susceptible individuals, negative cognitive styles and tendencies to ruminate can interact to create the conditions for the 'perfect storm': depression that is triggered by stressful events [86]. With continued practice of mindfulness, individuals may be better able to notice these patterns, allowing for disengagement from ruminative thought patterns (i.e. not being caught up in the thoughts and believing them as 'true') and selection of how to relate to these experiences, rather than 'automatically' reacting to them [16]. Indeed, studies have suggested that decreases in distress scores following MT are mediated by a decreased tendency to ruminate [87]. These findings suggest that MT may be effective in targeting habitual ruminative thinking, with consequent reduction in stress and depression [27].

Stress and ruminative thought patterns have also been linked to SUDs. For example, individuals' elevated rumination scores have been shown to predict substance abuse [69,88], and although efforts to avoid or suppress ruminative or 'unwanted' thoughts are used commonly in attempts to manage cravings and relapse [89], thought suppression has been shown to lead to stronger expectancies after cue exposure [90]. Interestingly, decreases in alcohol consumption following mindfulness-based treatment have been shown to be mediated partially by decreases in thought suppression indices such as avoidance [26]. These findings suggest that in dually diagnosed individuals, attentional focus on thoughts may be more effective than attempts to suppress them in decreasing their influence on behavior.

#### Neurobiological perspectives

It is hypothesized that specific areas of the medial PFC may be active during ruminative and/or wandering mindstates (dubbed the 'default mode') [91,92]. These regions have been found to play a role in linking subjective experiences through time [93,94], holding memory of traits of the self [95,96], reflected self-knowledge [97,98] and aspirations for the future [99]. Without this 'narrative self-reference', or sense of identity through time, stress reactivity and rumination would not be possible [100,101].

The mechanisms by which MT influences default mode functioning are being explored. In a recent study in which participants were instructed to either elaborate on current cognitions (narrative focus: NF) or to attend to somatic sensations and merely note any cognitions without elaborating on them (experiential focus: EF), investigators found midline cortical activation during NF compared to EF [102]. After MT, midline cortical structures showed decreased activity in EF versus NF [102]. One interpretation of these data is that momentary selfexperience may provide a non-self-related cortical 'task' which may suppress midline cortical activity. These findings are corroborated by studies showing increased gamma-band oscillation in the brains of long-term meditators (suggesting increased neuronal synchronization [103]) and behavioral studies showing improved attentional regulation with meditation training [104–106]. Together, these data suggest that present-centered attentional focus not only moves the individual away from the habitual default-mode thought process, but also manifests behaviorally. From these data, one might hypothesize that MT would benefit dually diagnosed individuals through improved attentional focus, with consequent reduction in stress-induced ruminative thought patterns, as well as more rapid recognition of these once they have been engaged. These would probably be reflected in 'lateralization' of brain activation patterns, as seen by Farb and colleagues in healthy individuals. Future studies in this population using neuroimaging attentional tasks will be informative in testing these hypotheses.

#### Acceptance

#### Theoretical perspectives

A second major component of mindfulness, acceptance, involves a non-judging/non-attached view of experiences. This perspective leads to an understanding of thoughts and sensations as 'transient mental events' rather than as reality or a reflection of the self. As with attentional focus, non-attached observation, or metacognition, has been hypothesized to reduce perpetuation of harmful thought patterns by shifting perspective of uncomfortable or unpleasant thoughts from 'real' or 'true' to viewing them as passing mental events. It should be noted that although acceptance is reported to be beneficial in MT as well as therapies that incorporate mindfulness [107,108], it is unclear whether a non-attached viewpoint fosters acceptance, acceptance fosters a nonattached viewpoint, or whether the effects are bidirectional, and to what degree these are an extension or result of insights gained from the practice of attention.

#### Clinical perspectives

The utility of a non-attached mode of experience in individuals with depression and/or addictions is gaining increasing support. Negative affect has been shown to predict relapse to both cigarettes and drug use [109– 111]. Also, higher intensity of negative affect has been correlated with longer duration of use [112], and worse distress tolerance has been associated with decreased abstinence [113]. Acceptance of distressing thoughts lessens reactivity and decreases attempts to avoid or suppress experiences, which have been linked to worsened outcomes [80,114]. For example, in clinical populations, MT has been associated with increases in the ability to 'let go of' (i.e. disengage from) negative automatic thoughts and decreases in the tendency towards negative automatic thinking [115]. Individuals with SUDs often report experiencing 'urges' to use substances, and many state that the experience of an urge is increasingly tolerated and managed when working with a mindfulness approach [24,116-118]. Additionally, decreases in substance use following MT have been shown to be mediated partially by decreased avoidance, but not frequency, of intrusive thoughts [26]. Further, distress tolerance, drawing from acceptance and commitment therapybased approaches, has shown preliminary utility in smoking cessation [119]. Finally, alcohol- and cocainedependent individuals have shown attenuation of selfreported anxiety and drug cravings during stress, with concomitant adaptive shifts in autonomic nervous system function, while remaining fully engaged with their experiences [24]. One might expect associated normalization in plasma cortisol concentrations, as has been suggested by studies of individuals with heart disease and cancer who have undergone MT [120,121], although these studies are yet to be reported. Together, these suggest that through acceptance of both mood- and drug-related ruminative thought patterns, MT may show increased efficacy in individuals with comorbid MDD and SUDs, where other treatments fail.

#### Neurobiological perspectives

Meta-cognitive skills, such as inhibition of secondary elaborative processing, may be fostered by MT, because attentional capacity is not being consumed by elaborative thinking [15,104]. Supporting this, Farb and colleagues have shown a decoupling of insula-vmPFC activity and an increased coupling of the insula with dlPFC after MT [102]. Further, meditation practice has been associated with thickened right insular and somatosensory cortices [122,123]. These data suggest a movement away from self-referential experiences (related to midline PFC activation) towards those that are more objectively observed/ felt and easily accepted. As stress has been shown to be associated with reduced dlPFC activation during particular tasks, with concomitant increased activation of the default-mode network, this lateralized brain activation pattern may also signal more adaptive responses with MT: neural resources are reallocated away from selfreferential, elaborative thinking towards task-specific responses, such as more accurate assessment of internal/ external situational stimuli and resultant skillful responses [124]. Consistent with this hypothesis, a recent study using the Stroop color-word interference task (which may predict treatment outcomes in addicted

individuals [47]) showed that MT was associated with improved cognitive flexibility [125]. We would hypothesize that dually diagnosed individuals would show similar adaptive patterns after MT with concomitant increased insular and dlPFC activation patterns, resulting in decreased perceived stress, drug use and depression severity.

# CONCLUSIONS AND FUTURE DIRECTIONS

Mindfulness training has shown promise in the treatment of both SUDs and MDD. Examination of the common neurobiological and behavioral dysfunction in these disorders suggests the promise of MT for dually diagnosed individuals. MT may help those with dual diagnosis decrease avoidance, tolerate unpleasant withdrawal and emotional states (stress-related), and unlearn maladaptive behaviors (rumination). Additionally, it may lessen the interactions between these processes, thus weakening their additive effects on depression and substance use.

We can now ask: do the commonalities in regional brain dysfunction between MDD and SUDs begin to approximate potential neural correlates of human suffering? If so, is this common to other psychiatric disorders, such as anxiety disorders, that share core features with both MDD and SUDs (e.g. unpleasant emotional states)? Would individuals with these disorders and comorbid SUDs also be helped by MT? Given the burgeoning research in discerning mechanisms of mindfulness and integrating MT in the treatment of psychiatric disorders, careful studies in dually diagnosed individuals have the potential to expand our knowledge of common pathophysiology greatly and provide effective treatments where few currently exist.

#### Declarations of interest

None.

## Acknowledgements

We would like to thank Bruce Rounsaville, Hedy Kober, Sharmin Ghaznavi, Zev Schuman-Olivier and Norman Farb for their helpful comments and discussions. This work was supported by funding from the following grants: NIDA K12-DA00167 (J.A.B.), T32-DA007238 (J.A.B.), R25 MH071584 (J.T.S.), R01 DA020908 (M.N.P.), P50 DA09241 (M.N.P.), NIAAA T32 AA 07455-24 (S.B.), and the VA VISN1 MIRECC (M.N.P.).

### References

1. Co-Occurring State Incentive Grants (COSIG) Clinical Protocol Committee. Definition of the Co-Occurring Disorders *Population*. Rockville, MD: Substance Abuse and Mental Health Services Administration (SAMHSA); 2006.

- 2. Kilgus M., Maxmen J. S., Ward N. G. Essential Psychopathology and Its Treatment. New York: Norton; 2009.
- Lopez A. D., Mathers C. D. Measuring the global burden of disease and epidemiological transitions: 2002–2030. *Ann Trop Med Parasitol* 2006; 100: 481–99.
- 4. Davis L. L., Rush J. A., Wisniewski S. R., Rice K., Cassano P., Jewell M. E. *et al.* Substance use disorder comorbidity in major depressive disorder: an exploratory analysis of the Sequenced Treatment Alternatives to Relieve Depression cohort. *Comp Psychiatry* 2005; **46**: 81.
- Davis L., Uezato A., Newell J. M., Frazier E. Major depression and comorbid substance use disorders. *Curr Opin Psychiatry* 2008; 21: 14–8.
- Compton W. M., Conway K. P., Stinson F. S., Grant B. F. Changes in the prevalence of major depression and comorbid substance use disorders in the United States between 1991 and 1992 and 2001–2002. *Am J Psychiatry* 2006; 163: 2141–7.
- Feldman G. Cognitive and behavioral therapies for depression: overview, new directions, and practical recommendations for dissemination. *Psychiatr Clin N Am* 2007; 30: 39.
- Marlatt G. A., Donovan D. M. Relapse Prevention: Maintenance Strategies in the Treatment of Addictive Behaviors, 2nd edn. New York: Guilford Press; 2005.
- Finney J. W., Moos R. H. Psychosocial treatments for alcohol use disorders. In: Nathan P. E., Gorman J. M., editors. *A Guide to Treatments That Work*. New York: Oxford University Press; 2002, p. 157–68.
- Carroll K. M., Onken L. S. Behavioral therapies for drug abuse. *Am J Psychiatry* 2005; 162: 1452–60.
- Quyen Q. T., Brent M. Treatments for patients with dual diagnosis: a review. *Alcohol Clin Exp Res* 2007; 31: 513– 36.
- Feldstein S. W., Ginsburg J. I. D. Motivational interviewing with dually diagnosed adolescents in juvenile justice settings. *Brief Treat Crisis Interv* 2006; 6: 218–33.
- Carroll K. M. Behavioral therapies for co-occurring substance use and mood disorders. *Biol Psychiatry* 2004; 56: 778.
- 14. Brown S. A., Glasner-Edwards S. V., Tate S. R., McQuaid J. R., Chalekian J., Granholm E. Integrated cognitive behavioral therapy versus twelve-step facilitation therapy for substance-dependent adults with depressive disorders. *J Psychoact Drugs* 2006; **38**: 449–60.
- Bishop S. R., Lau M., Shapiro S., Carlson L., Anderson N. D., Carmody J. et al. Mindfulness: a proposed operational definition. *Clin Psychol* 2004; 11: 230–41.
- Teasdale J. D., Segal Z., Williams J. M. G. How does cognitive therapy prevent depressive relapse and why should attentional control (mindfulness) training help? *Behav Res Ther* 1995; 33: 25.
- 17. Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: theoretical considerations and preliminary results. *Gen Hosp Psychiatry* 1982; 4: 33–47.
- Kabat-Zinn J., Lipworth L., Burney R. The clinical use of mindfulness meditation for the self-regulation of chronic pain. *J Behav Med* 1985; 8: 163–90.
- Roemer L., Orsillo S. M. Expanding our conceptualization of and treatment for generalized anxiety disorder: integrating mindfulness/acceptance-based approaches with

existing cognitive–behavioral models. *Clin Psychol Sci Pract* 2002; **9**: 54–68.

- Kabat-Zinn J., Massion A. O., Kristeller J., Peterson L. G., Fletcher K. E., Pbert L. *et al.* Effectiveness of a meditationbased stress reduction program in the treatment of anxiety disorders. *Am J Psychiatry* 1992; 149: 936–43.
- Miller J. J., Fletcher K., Kabat-Zinn J. Three-year follow-up and clinical implications of a mindfulness meditationbased stress reduction intervention in the treatment of anxiety disorders. *Gen Hosp Psychiatry* 1995; 17: 192– 200.
- Evans S., Ferrando S., Findler M., Stowell C., Smart C., Haglin D. Mindfulness-based cognitive therapy for generalized anxiety disorder. *J Anxiety Disord* 2008; 22: 716–21.
- 23. Zgierska A., Rabago D., Zuelsdorff M. *et al.* Mindfulness meditation for alcohol relapse prevention: a feasibility pilot study. *J Addict Med* 2008; **2**: 165–73.
- Brewer J. A., Sinha R., Chen J. A. *et al.* Mindfulness training and stress reactivity in substance abuse: results from a randomized, controlled stage 1 pilot study. *Subst Abuse* 2009; 30: 306–17.
- 25. Hoppes K. The application of mindfulness-based cognitive interventions in the treatment of co-occurring addictive and mood disorders. *CNS Spectr* 2006; 11: 829–51.
- Bowen S., Witkiewitz K., Dillworth T. M., Marlatt G. A. The role of thought suppression in the relationship between mindfulness meditation and alcohol use. *Addict Behav* 2007; 32: 2324–8.
- 27. Teasdale J. D., Segal Z. V., Williams J. M., Ridgeway V. A., Soulsby J. M., Lau M. A. Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. J Consult Clin Psychol 2000; 68: 615–23.
- Ma S. H., Teasdale J. D. Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *J Consult Clin Psychol* 2004; 72: 31–40.
- 29. Ospina M. B., Bond K., Karkhaneh M., Buscemi N., Dryden D. M., Barnes V. *et al.* Clinical trials of meditation practices in health care: characteristics and quality. *J Altern Complem Med* 2008; 14: 1199–213.
- Bowen S., Witkiewitz K., Dillworth T. M., Chawla N., Simpson T. L., Ostafin B. D. *et al.* Mindfulness meditation and substance use in an incarcerated population. *Psychol Addict Behav* 2006; 20: 343–7.
- Bowen S., Chawla N., Collins S. E., Witkiewitz K., Hsu S., Grow J. *et al.* Mindfulness-based relapse prevention for substance use disorders: a pilot efficacy trial. *Subst Abuse* 2009; **30**: 295–305.
- Kendler K. S., Gardner C. O., Prescott C. A. Clinical characteristics of major depression that predict risk of depression in relatives. *Arch Gen Psychiatry* 1999; 56: 322–7.
- 33. Sanacora G., Gueorguieva R., Epperson C. N., Wu Y.-T., Appel M., Rothman D. L. *et al.* Subtype-specific alterations of {gamma}-aminobutyric acid and glutamate in patients with major depression. *Arch Gen Psychiatry* 2004; 61: 705–13.
- Belmaker R. H., Agam G. Major depressive disorder. N Engl J Med 2008; 358: 55–68.
- 35. Caspi A., Sugden K., Moffitt T. E., Taylor A., Craig I. W., Harrington H. *et al.* Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 2003; **301**: 386–9.
- 36. Brody A. L., Saxena S., Mandelkern M. A., Fairbanks L. A.,

Ho M. L., Baxter L. R. Brain metabolic changes associated with symptom factor improvement in major depressive disorder. *Biol Psychiatry* 2001; **50**: 171.

- 37. Drevets W. C., Price J. L., Furey M. L. Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. *Brain Struct Funct* 2008; **213**: 93–118.
- 38. Goldapple K., Segal Z., Garson C., Lau M., Bieling P., Kennedy S. *et al.* Modulation of cortical–limbic pathways in major depression: treatment-specific effects of cognitive behavior therapy. *Arch Gen Psychiatry* 2004; 61: 34–41.
- Mayberg H. S., Lozano A. M., Voon V., McNeely H. E., Seminowicz D., Hamani C. *et al.* Deep brain stimulation for treatment-resistant depression. *Neuron* 2005; 45: 651– 60.
- 40. London E. D., Ernst M., Grant S., Bonson K., Weinstein A. Orbitofrontal cortex and human drug abuse: functional imaging. *Cereb Cortex* 2000; **10**: 334–42.
- Everitt B. J., Robbins T. W. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* 2005; 8: 1481–9.
- Wilson S. J., Sayette M. A., Fiez J. A. Prefrontal responses to drug cues: a neurocognitive analysis. *Nat Neurosci* 2004; 7: 211.
- 43. Goldstein R. Z., Tomasi D., Rajaram S., Cottone L. A., Zhang L., Maloney T. *et al.* Role of the anterior cingulate and medial orbitofrontal cortex in processing drug cues in cocaine addiction. *Neuroscience* 2007; **144**: 1153–9.
- 44. Olausson P., Jentsch J. D., Krueger D. D., Tronson N. C., Nairn A. C., Taylor J. R. Orbitofrontal cortex and cognitive–motivational impairments in psychostimulant addiction: evidence from experiments in the non-human primate. *Ann NY Acad Sci* 2007; **1121**: 610–38.
- 45. Bolla K. I., Eldreth D. A., London E. D., Kiehl K. A., Mouratidis M., Contoreggi C. *et al.* Orbitofrontal cortex dysfunction in abstinent cocaine abusers performing a decision-making task. *NeuroImage* 2003; 19: 1085.
- 46. Hester R., Garavan H. Executive dysfunction in cocaine addiction: evidence for discordant frontal, cingulate, and cerebellar activity. *J Neurosci* 2004; **24**: 11017–22.
- Brewer J. A., Worhunsky P. D., Carroll K. M., Rounsaville B. J., Potenza M. N. Pretreatment brain activation during stroop task is associated with outcomes in cocainedependent patients. *Biol Psychiatry* 2008; 64: 998– 1004.
- Barraclough D. J., Conroy M. L., Lee D. Prefrontal cortex and decision making in a mixed-strategy game. *Nat Neurosci* 2004; 7: 404–10.
- 49. MacDonald A. W., Cohen J. D., Stenger V. A., Carter C. S. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science* 2000; **288**: 1835–8.
- Mansouri F. A., Buckley M. J., Tanaka K. Mnemonic function of the dorsolateral prefrontal cortex in conflictinduced behavioral adjustment. *Science* 2007; 318: 987– 90.
- Phelps E. A., LeDoux J. E. Contributions of the amygdala to emotion processing: from animal models to human behavior. *Neuron* 2005; 48: 175–87.
- 52. Gottfried J. A., O'Doherty J., Dolan R. J. Encoding predictive reward value in human amygdala and orbitofrontal cortex. *Science* 2003; **301**: 1104–7.
- 53. Fanselow M. S., Gale G. D. The amygdala, fear, and memory. *Ann NY Acad Sci* 2003; **985**: 125–34.

- Damasio A. R., Grabowski T. J., Bechara A., Damasio H., Ponto L. L. B., Parvizi J. *et al.* Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nat Neurosci* 2000; 3: 1049–56.
- 55. Paus T., Barrett J. Transcranial magnetic stimulation (TMS) of the human frontal cortex: implications for repetitive TMS treatment of depression. J Psychiatry Neurosci Biobehav Rev 2004: 29: 268–79.
- Childress A. R., Mozley P. D., McElgin W., Fitzgerald J., Reivich M., O'Brien C. P. Limbic activation during cue-induced cocaine craving. *Am J Psychiatry* 1999; 156: 11–8.
- 57. Fuchs R. A., Evans K. A., Parker M. P., See R. E. Differential involvement of orbitofrontal cortex subregions in conditioned cue-induced and cocaine-primed reinstatement of cocaine seeking in rats. *J Neurosci* 2004; 24: 6600–10.
- 58. Brody A. L., Mandelkern M. A., London E. D., Childress A. R., Lee G. S., Bota R. G. *et al.* Brain metabolic changes during cigarette craving. *Arch Gen Psychiatry* 2002; 59: 1162–72.
- 59. Wang Z., Faith M., Patterson F., Tang K., Kerrin K., Wileyto E. P. *et al.* Neural substrates of abstinence-induced cigarette cravings in chronic smokers. *J Neurosci* 2007; 27: 14035–40.
- Garavan H., Pankiewicz J., Bloom A., Cho J. K., Sperry L., Ross T. J. *et al.* Cue-induced cocaine craving: neuroanatomical specificity for drug users and drug stimuli. *Am J Psychiatry* 2000; **157**: 1789–98.
- Naqvi N. H., Rudrauf D., Damasio H., Bechara A. Damage to the insula disrupts addiction to cigarette smoking. *Science* 2007; 315: 531–4.
- Paulus M. P., Tapert S. F., Schuckit M. A. Neural activation patterns of methamphetamine-dependent subjects during decision making predict relapse. *Arch Gen Psychiatry* 2005; 62: 761–8.
- 63. Nolen-Hoeksema S., Wisco B. E., Lyubomirsky S. Rethinking rumination. *Perspect Psychol Sci* 2008; **3**: 400–24.
- 64. Lyubomirsky S., Caldwell N. D., Nolen-Hoeksema S. Effects of ruminative and distracting responses to depressed mood on retrieval of autobiographical memories. *J Pers Soc Psychol* 1998; **75**: 166–77.
- 65. Davis R. N., Nolen-Hoeksema S. Cognitive inflexibility among ruminators and nonruminators. *Cogn Ther Res* 2000; **24**: 699.
- 66. Joorman J., Gotlib I. H. Inhibitory dysfunction, rumination, and mood regulation in depression, Paper presented at the annual meeting of the Association for the Advancement of Behavior Therapy, Washington, DC; 2005.
- 67. Ray R. D., Ochsner K. N., Cooper J. C., Robertson E. R., Gabrieli J. D. E., Gross J. J. Individual differences in trait rumination and the neural systems supporting cognitive reappraisal. *Cogn Affect Behav Neurosci* 2005; 5: 156–68.
- 68. Siegle G. J., Steinhauer S. R., Thase M. E., Stenger V. A., Carter C. S. Can't shake that feeling: event-related fMRI assessment of sustained amygdala activity in response to emotional information in depressed individuals. *Biol Psychiatry* 2002; **51**: 693.
- 69. Nolen-Hoeksema S., Harrell Z. A. Rumination, depression, and alcohol use: tests of gender differences. *J Cogn Psychother* 2002; **16**: 391–403.
- 70. Sinha R. How does stress increase risk of drug abuse and relapse? *Psychopharmacology (Berl)* 2001; **158**: 343–59.
- 71. Kessler R. C. The effects of stressful life events on depression. *Ann Rev Psychol* 1997; **48**: 191–214.

- Burke H. M., Davis M. C., Otte C., Mohr D. C. Depression and cortisol responses to psychological stress: a metaanalysis. *Psychoneuroendocrinology* 2005; 30: 846.
- 73. Kosten T. A., Zhang X. Y., Kehoe P. Chronic neonatal isolation stress enhances cocaine-induced increases in ventral striatal dopamine levels in rat pups. *Brain Res Dev Brain Res* 2003; 141: 109–16.
- Yavich L., Tiihonen J. Ethanol modulates evoked dopamine release in mouse nucleus accumbens: dependence on social stress and dose. *Eur J Pharmacol* 2000; 401: 365– 73.
- Sinha R., Garcia M., Paliwal P., Kreek M. J., Rounsaville B. J. Stress-induced cocaine craving and hypothalamic– pituitary–adrenal responses are predictive of cocaine relapse outcomes. *Arch Gen Psychiatry* 2006; 63: 324– 31.
- Bradley B. P., Phillips G., Green L., Gossop M. Circumstances surrounding the initial lapse to opiate use following detoxification. *Br J Psychiatry* 1989; 154: 354–9.
- Kalivas P. W., Duffy P. Similar effects of daily cocaine and stress on mesocorticolimbic dopamine neurotransmission in the rat. *Biol Psychiatry* 1989; 25: 913–28.
- Nash J. F. Jr, Maickel R. P. The role of the hypothalamic– pituitary–adrenocortical axis in post-stress induced ethanol consumption by rats. *Prog Neuropsychopharmacol Biol Psychiatry* 1988; 12: 653–71.
- 79. Oswald L. M., Wong D. F., McCaul M., Zhou Y., Kuwabara H., Choi L. *et al.* Relationships among ventral striatal dopamine release, cortisol secretion, and subjective responses to amphetamine. *Neuropsychopharmacology* 2005; **30**: 821–32.
- Leary M. R., Adams C. E., Tate E. B. Hypo-egoic selfregulation: exercising self-control by diminishing the influence of the self. *J Pers* 2006; 74: 1803–32.
- Allen N. B., Chambers R., Knight W. Mindfulness-based psychotherapies: a review of conceptual foundations, empirical evidence and practical considerations. *Aust NZ J Psychiatry* 2006; 40: 285–94.
- Brown K. W., Ryan R. M. The benefits of being present: mindfulness and its role in psychological well-being. *J Pers Soc Psychol* 2003; 84: 822–48.
- Lutz A., Slagter H. A., Dunne J. D., Davidson R. J. Attention regulation and monitoring in meditation. *Trends Cogn Sci* 2008; 12: 163–9.
- Breslin F. C., Zack M., McMain S. An informationprocessing analysis of mindfulness: implications for relapse prevention in the treatment of substance abuse. *Clin Psychol Sci Pract* 2002; 9: 275–99.
- Segal Z. V., Gemar M., Williams S. Differential cognitive response to a mood challenge following successful cognitive therapy or pharmacotherapy for unipolar depression. *J Abnorm Psychol* 1999; 108: 3–10.
- Robinson M. S., Alloy L. B. Negative cognitive styles and stress-reactive rumination interact to predict depression: a prospective study. *Cogn Ther Res* 2003; 27: 275–92.
- 87. Jain S., Shapiro S. L., Swanick S., Roesch S. C., Mills P. J., Bell I. *et al.* A randomized controlled trial of mindfulness meditation versus relaxation training: effects on distress, positive states of mind, rumination, and distraction. *Ann Behav Med* 2007; 33: 11–21.
- Nolen-Hoeksema S., Stice E., Wade E., Bohon C. Reciprocal relations between rumination and bulimic, substance abuse, and depressive symptoms in female adolescents. *J Abnorm Psychol* 2007; 116: 198–207.

- Salkovskis P. M., Reynolds M. Thought suppression and smoking cessation. *Behav Res Ther* 1994; 32: 193–201.
- Palfai T. P., Monti P. M., Colby S. M., Rohsenow D. J. Effects of suppressing the urge to drink on the accessibility of alcohol outcome expectancies. *Behav Res Ther* 1997; 35: 59–65.
- Gusnard D. A., Akbudak E., Gordon L. S., Raichle M. E. Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc Natl Acad Sci USA* 2001; 98: 4259.
- 92. Christoff K., Gordon A. M., Smallwood J., Smith R., Schooler J. W. Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proc Natl Acad Sci USA* 2009; 106: 8719–24.
- 93. Ulric N. The roots of self-knowledge: perceiving self, it, and thou. *Ann NY Acad Sci* 1997; **818**: 19–33.
- 94. Northoff G., Bermpohl F. Cortical midline structures and the self. *Trends Cogn Sci* 2004; 8: 102.
- 95. Macrae C. N., Moran J. M., Heatherton T. F., Banfield J. F., Kelley W. M. Medial prefrontal activity predicts memory for self. *Cereb Cortex* 2004; 14: 647–54.
- 96. Fossati P., Hevenor S. J., Graham S. J., Grady C., Keightley M. L., Craik F. *et al.* In search of the emotional self: an fMRI study using positive and negative emotional words. *Am J Psychiatry* 2003; **160**: 1938–45.
- Lieberman M. D., Jarcho J. M., Satpute A. B. Evidencebased and intuition-based self-knowledge: an fMRI study. *J Pers Soc Psychol* 2004; 87: 421–35.
- Ochsner K. N., Beer J. S., Robertson E. R., Cooper J. C., Gabrieli J. D. E., Kihsltrom J. F. *et al.* The neural correlates of direct and reflected self-knowledge. *NeuroImage* 2005; 28: 797.
- 99. Johnson M. K., Raye C. L., Mitchell K. J., Touryan S. R., Greene E. J., Nolen-Hoeksema S. Dissociating medial frontal and posterior cingulate activity during selfreflection. *Soc Cogn Affect Neurosci* 2006; 1: 56–64.
- 100. Mason M. F., Norton M. I., Van Horn J. D., Wegner D. M., Graffon S. T., Macrae C. N. Wandering minds: the default network and stimulus-independent thought. *Science* 2007; **315**: 393.
- Gallagher S. Philosophical conceptions of the self: implications for cognitive science. *Trends Cogn Sci* 2000; 4: 14.
- 102. Farb N. A. S., Segal Z. V., Mayberg H., Bean J., McKeon D., Fatima Z. *et al.* Attending to the present: mindfulness meditation reveals distinct neural modes of self-reference. *Soc Cogn Affect Neurosci* 2007; 2: 313–22.
- 103. Lutz A., Greischar L. L., Rawlings N. B., Ricard M., Davidson R. J. Long-term meditators self-induce high-amplitude gamma synchrony during mental practice. *Proc Natl Acad Sci USA* 2004; 101: 16369–73.
- 104. Jha A. P., Krompinger J., Baime M. J. Mindfulness training modifies subsystems of attention. *Cogn Affect Behav Neuro*sci 2007; 7: 109–19.
- 105. Srinivasan N., Baijal S. Concentrative meditation enhances preattentive processing: a mismatch negativity study. *Neuroreport* 2007; 18: 1709–12.
- 106. Brefczynski-Lewis J. A., Lutz A., Schaefer H. S., Levinson D. B., Davidson R. J. Neural correlates of attentional expertise in long-term meditation practitioners. *Proc Natl Acad Sci* 2007; 104: 11483–8.
- 107. Shearin E. N., Linehan M. M. Dialectical behavior therapy for borderline personality disorder: theoretical and

empirical foundations. *Acta Psychiatr Scand Suppl* 1994; **379**: 61–8.

- 108. Hayes S. C., Luoma J. B., Bond F. W., Masuda A., Lillis J. Acceptance and commitment therapy: model, processes and outcomes. *Behav Res Ther* 2006; **44**: 1–25.
- Marlatt G. A., Gordon J. R. Relapse Prevention: Maintenance Strategies in the Treatment of Addictive Behaviors. New York: Guilford Press; 1985.
- 110. Kenford S. L., Smith S. S., Wetter D. W., Jorenby D. E., Fiore M. C., Baker T. B. Predicting relapse back to smoking: contrasting affective and physical models of dependence. *J Consult Clin Psychol* 2002; **70**: 216–27.
- 111. Lowman C., Allen J., Stout R. L. Replication and extension of Marlatt's taxonomy of relapse precipitants: overview of procedures and results. The Relapse Research Group. *Addiction* 1996; 91: S51–71.
- Brown B. S. Relapse prevention in substance misuse: introduction. Int J Addict 1990; 25: 1081–3.
- 113. Daughters S. B., Lejuez C. W., Kahler C. W., Strong D. R., Brown R. A. Psychological distress tolerance and duration of most recent abstinence attempt among residential treatment-seeking substance abusers. *Psychol Addict Behav* 2005; 19: 208–11.
- 114. Wilson K. G., Murrell A. R. Values work in acceptance and commitment therapy: setting a course for behavioral treatment. In: Hayes S. C., Follette V., Linehan M. M., editors. *Mindfulness and Acceptance: Expanding the Cognitive-Behavioral Tradition*. New York: Guilford Press; 2004, p. 120–51.
- 115. Frewen P. A., Evans E. M., Maraj N., Dozois D. J. A., Partridge K. Letting go: mindfulness and negative automatic thinking. *Cogn Ther Res* **32**: 758–74.
- 116. Bowen S., Marlatt G. A. Surfing the urge: brief mindfulness-based intervention for college student smokers. *Psychol Addict Behav* 2009; **23**: 666–71.

- 117. Linehan M. Cognitive-Behavioral Treatment of Personality Disorder. New York: Guilford Press; 1993.
- 118. Marlatt G. A. Buddhist philosophy and the treatment of addictive behavior. *Cogn Behav Prac* 2002; 9: 44–50.
- 119. Brown R. A., Palm K. M., Strong D. R., Lejuez C. W., Kahler C. W., Zvolensky M. J. *et al.* Distress tolerance treatment for early-lapse smokers: rationale, program description, and preliminary findings. *Behav Modif* 2008; 32: 302–32.
- 120. Carlson L. E., Speca M., Patel K. D., Goodey E. Mindfulnessbased stress reduction in relation to quality of life, mood, symptoms of stress and levels of cortisol, dehydroepiandrosterone sulfate (DHEAS) and melatonin in breast and prostate cancer outpatients. *Psychoneuroendocrinology* 2004; **29**: 448–74.
- 121. Robert McComb J. J., Tacon A., Randolph P., Caldera Y. A pilot study to examine the effects of a mindfulness-based stress-reduction and relaxation program on levels of stress hormones, physical functioning, and submaximal exercise responses. *J Altern Complem Med* 2004; **10**: 819–27.
- 122. Holzel B. K., Ott U., Gard T., Hempel H., Weygandt M., Morgen K. *et al.* Investigation of mindfulness meditation practitioners with voxel-based morphometry. *Soc Cogn Affect Neurosci* 2008; **3**: 55–61.
- 123. Lazar S. W., Kerr C. E., Wasserman R. H., Gray J. R., Greve D. N., Treadway M. T. *et al.* Meditation experience is associated with increased cortical thickness. *Neuroreport* 2005; 16: 1893–7.
- 124. Qin S., Hermans E. J., Marle H. J. F., Luo J., Fernandez G. Acute psychological stress reduces working memoryrelated activity in the dorsolateral prefrontal cortex. *Biol Psychiatry* 2009; 66: 25–32.
- 125. Moore A., Malinowski P. Meditation, mindfulness and cognitive flexibility. *Conscious Cogn* 2009; **18**: 176–86.