

BRIEF REPORT

An Invariant Dimensional Liability Model of Gender Differences in Mental Disorder Prevalence: Evidence From a National Sample

Nicholas R. Eaton
University of Minnesota

Katherine M. Keyes
Columbia University

Robert F. Krueger
University of Minnesota

Steve Balsis
Texas A&M University

Andrew E. Skodol
Columbia University and University of Arizona

Kristian E. Markon
University of Iowa

Bridget F. Grant
National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD

Deborah S. Hasin
Columbia University

Epidemiological studies of categorical mental disorders consistently report that gender differences exist in many disorder prevalence rates and that disorders are often comorbid. Can a dimensional multivariate liability model be developed to clarify how gender impacts diverse, comorbid mental disorders? We pursued this possibility in the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC; $N = 43,093$). Gender differences in prevalence were systematic such that women showed higher rates of mood and anxiety disorders, and men showed higher rates of antisocial personality and substance use disorders. We next investigated patterns of disorder comorbidity and found that a dimensional internalizing-externalizing liability model fit the data well, where internalizing is characterized by mood and anxiety disorders, and externalizing is characterized by antisocial personality and substance use disorders. This model was gender invariant, indicating that observed gender differences in prevalence rates originate from women and men's different average standings on latent internalizing and externalizing liability dimensions. As hypothesized, women showed a higher mean level of internalizing, while men showed a higher mean level of externalizing. We discuss implications of these findings for understanding gender differences in psychopathology and for classification and intervention.

Keywords: comorbidity, gender differences, internalizing–externalizing, prevalence rates

Previous epidemiological studies have shown that there are sizable gender differences in the prevalence rates of many common mental disorders (for recent reviews, see Grant & Weissman, 2007; Shear, Halmi, Widiger, & Boyce, 2007; Widiger, 2007). For example, 12-month and lifetime prevalence rates from the National Comorbidity Survey indicated that women showed markedly higher (and often approximately double) prevalence rates of major

depression, dysthymia, generalized anxiety disorder, panic disorder, social phobia, and specific phobia than did men. In contrast, men showed higher prevalence rates of antisocial personality disorder and alcohol and drug dependence (Kessler, McGonagle, Swartz, Blazer, & Nelson, 1993; Kessler et al., 1994). Similar gender differences have been observed in the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC), the

This article was published Online First August 15, 2011.

Nicholas R. Eaton and Robert F. Krueger, Department of Psychology, University of Minnesota; Katherine M. Keyes and Deborah S. Hasin, Departments of Epidemiology and Psychiatry, Columbia University; Steve Balsis, Department of Psychology, Texas A&M University; Andrew E. Skodol, College of Physicians and Surgeons, Columbia University and Department of Psychiatry, University of Arizona; Kristian E. Markon, Department of Psychology, University of Iowa; Bridget F. Grant, Laboratory of Epidemiology and Biometry, Division of Intramural Clinical and Biological Research, National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD.

This research was supported by the following grants: U01AA018111, R01DA018652, and K05AA014223 (Hasin); and F31DA026689 (Keyes). The National Epidemiologic Survey on Alcohol and Related Conditions (NESARC) was sponsored by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) and funded, in part, by the Intramural Program of the NIAAA, with additional support from the National Institute on Drug Abuse.

Correspondence concerning this article should be addressed to Nicholas R. Eaton, Department of Psychology, University of Minnesota, 75 East River Road, Minneapolis, MN 55455. E-mail: nreaton@gmail.com

largest epidemiological study of psychopathology yet undertaken (Dawson, Goldstein, Moss, Li, & Grant, 2010; Keyes, Grant, & Hasin, 2008; Grant et al., 2004; Grant & Weissman, 2007; Trull, Jahng, Tomko, Wood, & Sher, 2010; Vesga-López et al., 2008).

The origins of these gender differences in prevalence rates are not well understood although various theories have been posited to explain how they arise. These explanations include response bias, differential service utilization rates, and various biological, social, and demographic influences (see Klose & Jacobi, 2004; Piccinelli & Wilkinson, 2000). Psychological explanations, such as increased rumination in women partially accounting for higher rates of unipolar depression, have also been posited (Nolen-Hoeksema, 1987; Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008).

These theories of gender differences focus primarily on specific disorders and rarely take comorbidity into account. A compelling account of mental disorder comorbidity focuses on unifying latent dimensional liabilities to experience multiple internalizing (mood and anxiety) or externalizing (antisocial and substance use) disorders (Eaton, South, & Krueger, 2010; Krueger, 1999; Slade & Watson, 2006; Vollebergh et al., 2001). Indeed, this internalizing–externalizing liability model is likely to frame key parts of the *metastructure*, or overall organization, of the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM–5)*; e.g., Andrews et al., 2009; Regier, Narrow, Kuhl, & Kupfer, 2011). The internalizing dimension can be bifurcated into distress and fear subfactors; distress relates to disorders such as major depression, dysthymia, and generalized anxiety, and fear relates to disorders such as panic disorder, social phobia, and specific phobia. The externalizing dimension is associated with disorders such as antisocial personality disorder and alcohol, nicotine, and drug dependence. Further, these factors relate to normal personality: Internalizing correlates with neuroticism/negative affectivity (Griffith et al., 2010), and externalizing correlates with disinhibition (Krueger et al., 2002).

When gender differences in prevalence rates and the internalizing–externalizing liability structure of psychopathology are considered simultaneously, the possibility of a unifying model of gender and comorbidity emerges. Specifically, women show significantly higher prevalence rates of internalizing disorders, while men show significantly higher rates of externalizing disorders (Grant & Weissman, 2007; Kessler et al., 1993, 1994). This observation suggests that gender differences in *categorical* prevalence rates might be due to gender differences in latent internalizing and externalizing liability *dimensions*. The utility of a dimensional liability model for public health, epidemiology, psychopathology, and intervention research would be notably enhanced if it could encompass the role of gender in mental disorder prevalence.

A few studies have evaluated the structure of psychopathology separately in women and men (e.g., Krueger, 1999; Kendler, Prescott, Myers, & Neale, 2003). However, we are aware of only two studies that have *formally tested* whether the latent structure of common mental disorders is gender invariant (Hicks et al., 2007; Kramer et al., 2008). While these studies were generally supportive of a gender invariant model, their findings' generalizability was limited by nonrepresentative samples. Further, neither study focused on diagnoses from the fourth edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM–IV)*; American Psychiatric Association, 1994). Addressing these limitations is critical,

given the potential use of a dimensional liability model to frame key aspects of *DSM–5*. If the model is to be applied to *DSM–5* mental disorders—and to both women and men—factorial invariance (a lack of bias) across gender should be demonstrated in a maximally representative sample.

The current study examined a nationally representative sample of 43,093 individuals assessed in the first wave of the NESARC. Our initial goal was to examine and present any gender differences in prevalence rates of common mental disorders in this sample. We were interested not only in the *significance* of these gender differences but also their *directionality*: We hypothesized that women would show significantly higher rates of internalizing disorders than men, and men would show significantly higher rates of externalizing disorders. Second, we sought to determine the latent comorbidity structure of these disorders in women and men, separately. Third, we aimed to formally test whether or not the emergent structures in women and men could be considered gender invariant (i.e., women and men showing equivalent structures of psychopathology). Finally, if invariance were found, we hypothesized that women would have a higher mean standing on the internalizing liability dimension than men, and men would have a higher mean standing on the externalizing liability dimension. The presence of a gender invariant structure of common mental disorders would indicate that these gender differences in latent internalizing and externalizing liabilities account for the observed gender differences in prevalence rates. That is, gender differences in the prevalence of different manifest categorical disorders would be a function of mean-level gender differences in underlying liability dimensions. As such, these underlying *dimensions*, as opposed to their manifestations as specific observed *categories* of psychopathology, would be highlighted as important organizing constructs for official nosologies and for research on the role of gender in psychopathology.

Method

Participants

This study utilized data from 43,093 individuals who participated in the first wave of the NESARC, conducted in 2001–2002. The NESARC study's design has been detailed elsewhere (Grant & Dawson, 2006). The first wave of NESARC was a representative sample of the civilian, noninstitutionalized United States population, aged 18 and older. Young adults, African Americans, and Hispanics were oversampled. Women composed 57% ($n = 24,575$). Race/ethnicity was selected by participants, using census-defined categories: White (56.9%), Hispanic or Latino (19.3%), African American (19.1%), Asian/Native Hawaiian/Pacific Islander (3.1%), and American Indian/Alaska Native (1.6%). Participants provided written informed consent after reviewing a complete description of the study.

Assessment

Lifetime and past-12-month *DSM–IV* diagnoses were made using the Alcohol Use Disorder and Associated Disabilities Interview Schedule—*DSM–IV* Version (AUDADIS–IV; Grant et al., 1995), a structured interview designed for experienced lay interviewers. Major depressive disorder, dysthymic disorder, general-

ized anxiety disorder, panic disorder, social phobia, specific phobia, alcohol dependence, nicotine dependence, marijuana dependence, other drug dependence, and antisocial personality disorder AUDADIS-IV diagnoses were examined. The other drug dependence variable was created to collapse relatively uncommon forms of drug dependence (i.e., stimulants, opioids, sedatives, tranquilizers, cocaine, solvents, hallucinogens, heroin, and any other drug not assessed) into one variable whose variance would be sufficient for covariance structure modeling; the internal consistency of this variable was good ($\alpha = .77$). In keeping with *DSM-IV* notions of personality disorder stability, antisocial personality disorder was assessed on a lifetime basis only; this lifetime diagnosis was used in both lifetime and 12-month analyses. The reliability of the AUDADIS-IV diagnoses examined have been reported elsewhere and are generally good to excellent (e.g., kappas = .42 to .84; see Hasin et al., 2005). Test-retest estimates for AUDADIS-IV disorders are similar to other structured interviews (e.g., the Diagnostic Interview Schedule [DIS], the Composite International Diagnostic Interview [CIDI]) used in large psychiatric epidemiologic surveys (reviewed in Wittchen, 1994). Further, the AUDADIS-IV has advantages over structured interviews such as the DIS, including assessment of clinically significant distress and impairment after the syndrome is fully characterized (Hasin et al., 2005).

Statistical Analyses

All analyses were conducted in Mplus version 6 (Muthén & Muthén, 2010) using the Mplus defaults of delta parameterization and WLSMV estimator. WLSMV allowed us to treat diagnostic variables as categorical and use the NESARC's weighting, clustering, and stratification variables. Odds ratios used men as the reference comparison group. To evaluate model fit in confirmatory factor analyses (CFAs), we considered the comparative fit index (CFI), Tucker-Lewis index (TLI), RMSEA, and the number of freely estimated parameters in the model. CFI/TLI values $> .95$ and RMSEA values $< .06$ suggest good model fit (Hu & Bentler, 1999). Based on simulations, Cheung and Rensvold (2002) proposed that a CFI difference critical value of .01 be used to test whether the addition of constraints leads to notably worse model fit in factorial invariance studies. More

parsimonious models use fewer freely estimated parameters. In model comparisons, therefore, we defined the optimal model by means of the best fit (CFI, TLI, and RMSEA), model parsimony (number of free parameters), and the CFI critical difference of .01. Distress and fear loadings on higher-order internalizing were constrained to equality to ensure model identification.

Results

Prevalence Rates

Table 1 presents the prevalence rates for the disorders included in the current study, separately for men and women and for lifetime and 12-month disorders. All odds ratios were significant (all p s $< .001$ except other drug dependence at $p = .005$). Across lifetime and 12-month prevalence rates, women showed higher rates for all internalizing disorders, and men showed higher rates for all externalizing disorders.

Comorbidity Structure

In CFAs (see Table 2), guided by previous studies and exploratory factor analyses (not reported here for brevity), we parameterized each diagnosis to load on one of three factors: (1) distress: major depression, dysthymia, and generalized anxiety disorder; (2) fear: panic disorder, social phobia, and specific phobia; and (3) externalizing: alcohol dependence, nicotine dependence, marijuana dependence, other drug dependence, other drug dependence, and antisocial personality disorder. Distress and fear were parameterized to load on a higher-order internalizing factor, which was allowed to correlate with the externalizing factor. This internalizing-externalizing model provided a very good fit in the total sample for both lifetime and 12-month diagnoses. Within each gender modeled separately, this internalizing-externalizing model continued to fit very well for lifetime and 12-month diagnoses.

Invariance

Because an internalizing-externalizing model fit well for both women and men, our next question was how similar these models

Table 1
Disorder Prevalence Rates and Odds Ratios by Gender

Disorder	Lifetime disorders			12-month disorders		
	Women (%)	Men (%)	Odds ratio	Women (%)	Men (%)	Odds ratio
Depression	22.9	13.1	1.46 (1.41–1.51)	10.1	5.5	1.38 (1.32–1.45)
Dysthymia	6.2	3.5	1.31 (1.25–1.38)	2.9	1.6	1.29 (1.20–1.39)
Generalized Anxiety	5.8	3.1	1.34 (1.27–1.42)	3.1	1.4	1.37 (1.28–1.47)
Panic Disorder	7.2	3.7	1.39 (1.32–1.47)	3.1	1.4	1.39 (1.29–1.49)
Social Phobia	5.8	4.3	1.16 (1.10–1.22)	3.4	2.1	1.22 (1.15–1.29)
Specific Phobia	12.4	6.2	1.47 (1.41–1.53)	9.6	4.6	1.46 (1.40–1.53)
Alcohol Dependence	8.0	17.4	0.63 (0.60–0.65)	2.3	5.4	0.68 (0.64–0.72)
Nicotine Dependence	15.6	20.0	0.84 (0.81–0.87)	11.5	14.1	0.88 (0.85–0.91)
Marijuana Dependence	0.9	1.7	0.77 (0.71–0.83)	0.2	0.5	0.72 (0.63–0.83)
Other Drug Dependence	1.4	2.2	0.84 (0.79–0.90)	0.3	0.5	0.83* (0.72–0.95)
Antisocial Personality	1.9	5.5	0.68 (0.59–0.66)			

Note. All odds ratios (OR) significant at $p < .001$, except * $p = .005$. Men are OR comparison group. 95% confidence intervals are given in parentheses. Antisocial personality disorder was only assessed as a lifetime disorder.

Table 2
Model Fit Statistics

Model	CFI	TLI	RMSEA	# Free
Total sample ($N = 43,093$)				
Lifetime diagnoses	.992	.989	.012	—
12-month diagnoses	.988	.984	.010	—
Women ($n = 24,575$)				
Lifetime diagnoses	.993	.991	.009	—
12-month diagnoses	.990	.987	.008	—
Men ($n = 18,518$)				
Lifetime diagnoses	.988	.984	.008	—
12-month diagnoses	.982	.976	.007	—
Multigroup (Women and Men)				
Lifetime diagnoses				
Unconstrained model	.991	.989	.012	48
Constrained model	.991	.989	.012	38
12-month diagnoses				
Unconstrained model	.987	.983	.010	48
Constrained model	.988	.986	.009	38

Note. Total sample analyses modeled women and men together. Multigroup analyses modeled women and men simultaneously as two separate groups. Unconstrained models allowed each gender to have unique model parameters; constrained (invariant) models constrained factor loadings and thresholds to equality across genders. CFI = comparative fit index; TLI = Tucker-Lewis index; RMSEA = root mean squared error of approximation; # Free = number of freely estimated parameters.

were across gender in terms of model parameters—that is, whether the magnitude of parameters differed by gender or whether they showed invariance. Tests of invariance for indicators such as diagnoses require methodology appropriate for modeling categorical variables (Millsap & Yun-Tein, 2004). In this approach, factor

loadings and thresholds are constrained to equality or freed, in tandem, across genders. In our first model (the “unconstrained model”), loadings and thresholds were free across genders, factor means were set to 0 in both genders, and scaling factors were fixed to 1 in both genders. In the second model (the “constrained model”), loadings and thresholds were constrained to equality across genders, factor means were set to 0 in men and were free in women, and scaling factors were fixed to 1 in men and were free in women. This model represented a gender invariant psychopathology structure.

We fit the unconstrained and constrained models in men and women simultaneously via a multiple group CFA, separately for lifetime and 12-month diagnoses (see Table 2). For lifetime diagnoses, the fits of the two models were identical, but the constrained model had fewer freely estimated parameters than the unconstrained model. The constrained model for lifetime diagnoses is depicted in Figure 1. For 12-month diagnoses, the constrained model had a better fit, with greater parsimony, than did the unconstrained model. For lifetime and 12-month diagnoses, the CFI critical difference of .01 was not exceeded, further supporting the constrained model. These findings indicated that, in addition to the general structure, factor loadings and thresholds for all diagnoses were equivalent for women and men. Thus, the structure of these common mental disorders, including the connections between individual diagnoses and the underlying factors, could be considered gender invariant.

In terms of factor means, means of these latent internalizing and externalizing factors were fixed to 0 in men and freely estimated as .445 and $-.378$ in women, respectively, for lifetime diagnoses and as .428 and $-.308$ for 12-month diagnoses. All mean gender differences were significant at $p < .01$. These standardized means




Figure 1. The constrained (gender invariant) model in women and men using lifetime diagnoses. Values are standardized factor loadings (all significant $p < .001$). Values before slash and bolded are for women; values after slash are for men. Values differ slightly across gender due to standardization. MDD = major depressive disorder; Dysth = dysthymic disorder; GAD = generalized anxiety disorder; Panic = panic disorder; Social = social phobia; Spec = specific phobia; ASPD = antisocial PD; Nic = nicotine dependence; Alc = alcohol dependence; Marij = marijuana dependence; Drug = other drug dependence. Arrows without numbers indicate unique variances, including error.

can be interpreted as z-scores (e.g., women were approximately .45 standard deviations *higher* on lifetime internalizing liability than men). Because complete factorial invariance had been established, these results demonstrated that the observed differences in the prevalence rates of the specific disorders modeled between women and men could be accounted for by the genders' different average levels of latent internalizing and externalizing.

Discussion

The current study sought to synthesize two lines of research—patterns of gender differences in prevalence rates and a potentially gender invariant latent structure of psychopathology—via factorial invariance analyses of liability dimensions underlying *DSM-IV* disorder comorbidity. We found that the underlying structure of common mental disorders was gender invariant, with significant gender differences in mean liability levels. This provides compelling evidence that observed gender differences in prevalence rates of many common mental disorders originate at the level of latent internalizing and externalizing liabilities.

Limitations

This study is not without its limitations. First, we examined lifetime diagnoses, which can be subject to memory biases. However, our results from lifetime diagnoses were highly congruent with results from 12-month diagnoses, which require much less retrospection. Second, our diagnostic information was collected by extensively trained lay interviewers rather than clinicians. This being said, it is noteworthy that the instrument used to assess symptomatology was fully structured, which resulted in generally good diagnostic reliability levels. Finally, the current study investigated only common mental disorders and, thus, did not include other debilitating forms of psychopathology, such as schizophrenia. There are indications that some symptoms of psychotic disorders may relate to a separate liability factor (e.g., thought disorder liability) while also showing associations with internalizing liability/neuroticism (e.g., Barrantes-Vidal, Ros-Morente, & Kwapil, 2009; Markon, 2010).

Implications

Classification. *DSM-IV* is currently under revision, and there has been a great deal of discussion about the general organization of *DSM-5* (Regier et al., 2011). Based largely on replications of the internalizing–externalizing model, an organizational meta-structure for many common disorders reflecting this structure has been advocated (Andrews et al., 2009). Our findings support this proposal in two ways. First, we replicated the internalizing–externalizing structure in the NESARC, the largest epidemiologic study of psychopathology yet undertaken. Second, our results indicated that this structure is gender invariant. The current study represents the first time that gender invariance has been tested and successfully incorporated into the internalizing–externalizing liability model of comorbidity among categorical *DSM-IV* mental disorders in a representative sample. Taken together, these findings support an internalizing–externalizing metastructure for many disorders in *DSM-5*, especially because the model is applicable in both women and men.

Gender differences research. Our conclusion—that the observed gender differences in prevalence rates systematically reflect gender differences in broad latent liability factors—ties together distinct lines of research and theory on gender differences in prevalence rates for specific disorders. For instance, one major theory to account for gender differences in depression involves the notion that women ruminate more frequently than men, focusing repetitively on their negative emotions and problems rather than engaging in more active problem solving (Nolen-Hoeksema, 1987; Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008). This theory can be readily extended to anxiety (and other internalizing disorders) by noting that neuroticism, or negative affectivity, is strongly related to rumination such that individuals who are more neurotic ruminate more frequently (Lam, Smith, Checkley, Rijdsdijk, & Sham, 2003). Neuroticism is also strongly related to ($r = .98$), and nearly isomorphic with, the latent internalizing dimension reflecting the multivariate comorbidity among *DSM-IV* mood and anxiety disorders (Griffith et al., 2010). This link between internalizing and trait neuroticism is itself accounted for largely by genetic effects (Hettema et al., 2006). Finally, previous research has indicated that women tend to report higher levels of trait neuroticism (as well as conscientiousness and agreeableness) on average than do men (e.g., Donnellan & Lucas, 2008), which mirrors our finding that women had significantly higher mean levels of internalizing than did men. It may be through neuroticism (and disinhibition-related traits in the case of externalizing and men; e.g., Krueger et al., 2002; Miller & Lynam, 2001; Slutske et al., 2002) that psychological processes impact latent propensities to experience comorbid mental disorders. Given that women tend to report higher frequencies of some stressful life events than men do prior to disorder onset (Harkness et al., 2010), the interaction between these liabilities and environmental stressors seems a particularly worthwhile focus for gender differences research.

Intervention and prevention. Our results support recent efforts to develop interventions that target latent disorder liabilities. For instance, both anxious and depressive symptoms often respond to the same pharmacologic interventions (Goldberg, Simms, Gater, & Krueger, 2011). Similarly, anxiety and depression both respond well to cognitive–behavioral therapy, and there have been efforts to develop psychotherapeutic interventions that address the shared internalizing liability rather than solely focusing on its manifestations (Barlow et al., 2011). Along these lines, prevention efforts that focus on gender-linked core psychological processes are likely to be effective in impacting multiple disorders. In women, these preventative measures might focus, for instance, on coping and cognitive restructuring skills to reduce the likelihood of rumination and cognitive distortions developing into clinically significant depression or anxiety. In men, prevention might focus on rewarding planful behaviors and shaping disinhibitory tendencies into outlets that are not destructive to the self or others.

References

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Andrews, G., Goldberg, D. P., Krueger, R. F., Carpenter, W. T., Jr., Hyman, S. E., Sachdev, P., & Pine, D. S. (2009). Exploring the feasibility of a meta-structure for DSM-V and ICD-11: Could it improve utility and validity? *Psychological Medicine: A Journal of Research in*

- Psychiatry and the Allied Sciences*, 39, 1993–2000. doi:10.1017/S0033291709990250
- Barlow, D. H., Farchione, T. J., Fairholme, C. P., Ellard, K. K., Boisseau, C. L., Allen, L. B., & Ehrenreich-May, J. (2011). *Unified protocol for transdiagnostic treatment of emotional disorders: Therapist guide*. New York: Oxford University Press.
- Barrantes-Vidal, N., Ros-Morente, A., & Kwapil, T. R. (2009). An examination of neuroticism as a moderating factor in the association of positive and negative schizotypy with psychopathology in a nonclinical sample. *Schizophrenia Research*, 115, 303–309. doi:10.1016/j.schres.2009.09.021
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling*, 9, 233–255. doi:10.1207/S15328007SEM0902_5
- Dawson, D. A., Goldstein, R. B., Moss, H. B., Li, T.-K., & Grant, B. F. (2010). Gender differences in the relationship of internalizing and externalizing psychopathology to alcohol dependence: Likelihood, expression and course. *Drug and Alcohol Dependence*, 112, 9–17. doi:10.1016/j.drugalcdep.2010.04.019
- Donnellan, M. B., & Lucas, R. E. (2008). Age differences in the Big Five across the life span: Evidence from two national samples. *Psychology and Aging*, 23, 558–566. doi:10.1037/a0012897
- Eaton, N. R., South, S. C., & Krueger, R. F. (2010). The meaning of comorbidity among common mental disorders. In T. Millon, R. F. Krueger, & E. Simonsen (Eds.), *Contemporary directions in psychopathology* (2nd ed., pp. 223–241). New York: Guilford Publications.
- Goldberg, D., Simms, L. J., Gater, R., & Krueger, R. F. (2011). Integration of dimensional spectra for depression and anxiety into categorical diagnoses for general medical practice. In D. A. Regier, W. E. Narrow, E. A. Kuhl, & D. J. Kupfer (Eds.), *The conceptual evolution of DSM-5* (pp. 19–35). Arlington, VA: American Psychiatric Publishing, Inc.
- Grant, B. F., & Dawson, D. A. (2006). Introduction to the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Research & Health*, 29, 74–78.
- Grant, B. F., Dawson, D. A., Stinson, F. S., Chou, S. P., Dufour, M. C., & Pickering, R. P. (2004). The 12-month prevalence and trends in DSM-IV alcohol abuse and dependence: United States, 1991–1992 and 2001–2002. *Drug and Alcohol Dependence*, 74, 223–234. doi:10.1016/j.drugalcdep.2004.02.004
- Grant, B. F., Harford, T. C., Dawson, D. A., Chou, P. S., & Pickering, R. P. (1995). The Alcohol Use Disorder and Associated Disabilities Interview Schedule (AUDADIS): Reliability of alcohol and drug modules in a general population sample. *Drug and Alcohol Dependence*, 39, 37–44. doi:10.1016/0376-8716(95)01134-K
- Grant, B. F., & Weissman, M. M. (2007). Gender and the prevalence of psychiatric disorders. In W. E. Narrow, M. B. First, P. J. Sirovatka, & D. A. Regier (Eds.), *Age and gender considerations in psychiatric diagnosis: A research agenda for DSM-V* (pp. 31–46). Washington, DC: American Psychiatric Association.
- Griffith, J. W., Zinbarg, R. E., Craske, M. G., Mineka, S., Rose, R. D., Waters, A. M., & Sutton, J. M. (2010). Neuroticism as a common dimension in the internalizing disorders. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*, 40, 1125–1136. doi:10.1017/S0033291709991449
- Harkness, K. L., Alavi, N., Monroe, S. M., Slavich, G. M., Gotlib, I. H., & Bagby, R. M. (2010). Gender differences in life events prior to onset of major depressive disorder: The moderating effect of age. *Journal of Abnormal Psychology*, 119, 791–803. doi:10.1037/a0020629
- Hasin, D. S., Goodwin, R. D., Stinson, F. S., & Grant, B. F. (2005). Epidemiology of major depressive disorder: Results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Archives of General Psychiatry*, 62, 1097–1106. doi:10.1001/archpsyc.62.10.1097
- Hettema, J. M., Neale, M. C., Myers, J. M., Prescott, C. A., & Kendler, K. S. (2006). A population-based twin study of the relationship between neuroticism and internalizing disorders. *American Journal of Psychiatry*, 163, 857–864. doi:10.1176/appi.ajp.163.5.857
- Hicks, B. M., Blonigen, D. M., Kramer, M. D., Krueger, R. F., Patrick, C. J., Iacono, W. G., & McGue, M. (2007). Gender differences and developmental change in externalizing disorders from late adolescence to early adulthood: A longitudinal twin study. *Journal of Abnormal Psychology*, 116, 433–447. doi:10.1037/0021-843X.116.3.433
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1–55. doi:10.1080/10705519909540118
- Kendler, K. S., Prescott, C. A., Myers, J., & Neale, M. C. (2003). The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Archives of General Psychiatry*, 60, 929–937. doi:10.1001/archpsyc.60.9.929
- Kessler, R. C., McGonagle, K. A., Swartz, M., Blazer, D. G., & Nelson, C. B. (1993). Sex and depression in the National Comorbidity Survey I: Lifetime prevalence, chronicity and recurrence. *Journal of Affective Disorders*, 29, 85–96. doi:10.1016/0165-0327(93)90026-G
- Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H.-U., & Kendler, K. S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. *Archives of General Psychiatry*, 51, 8–19.
- Keyes, K. M., Grant, B. F., & Hasin, D. S. (2008). Evidence for a closing gender gap in alcohol use, abuse, and dependence in the United States population. *Drug and Alcohol Dependence*, 93, 21–29. doi:10.1016/j.drugalcdep.2007.08.017
- Klose, M., & Jacobi, F. (2004). Can gender differences in the prevalence of mental disorders be explained by sociodemographic factors? *Archives of Women's Mental Health*, 7, 133–148. doi:10.1007/s00737-004-0047-7
- Kramer, M. D., Krueger, R. F., & Hicks, B. M. (2008). The role of internalizing and externalizing liability factors in accounting for gender differences in the prevalence of common psychopathological syndromes. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*, 38, 51–61. doi:10.1017/S0033291707001572
- Krueger, R. F. (1999). The structure of common mental disorders. *Archives of General Psychiatry*, 56, 921–926. doi:10.1001/archpsyc.56.10.921
- Krueger, R. F., Hicks, B. M., Patrick, C. J., Carlson, S. R., Iacono, W. G., & McGue, M. (2002). Etiologic connections among substance dependence, antisocial behavior, and personality: Modeling the externalizing spectrum. *Journal of Abnormal Psychology*, 111, 411–424. doi:10.1037/0021-843X.111.3.411
- Lam, D., Smith, N., Checkley, S., Rijdsdijk, F., & Sham, P. (2003). Effect of neuroticism, response style and information processing on depression severity in a clinically depressed sample. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*, 33, 469–479. doi:10.1017/S0033291702007304
- Markon, K. E. (2010). Modeling psychopathology structure: A symptom-level analysis of Axis I and II disorders. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*, 40, 273–288. doi:10.1017/S0033291709990183
- Miller, J. D., & Lynam, D. (2001). Structural models of personality and their relation to antisocial behavior: A meta-analytic review. *Criminology*, 39, 765–798. doi:10.1111/j.1745-9125.2001.tb00940.x
- Millsap, R. E., & Yun-Tein, J. (2004). Assessing factorial invariance in ordered-categorical measures. *Multivariate Behavioral Research*, 39, 479–515. doi:10.1207/S15327906MBR3903_4
- Muthén, L. K., & Muthén, B. O. (2010). *Mplus user's guide* (5th ed.). Los Angeles: Author.
- Nolen-Hoeksema, S. (1987). Sex differences in unipolar depression: Evidence and theory. *Psychological Bulletin*, 101, 259–282. doi:10.1037/0033-2909.101.2.259

- Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspectives on Psychological Science*, 3, 400–424. doi:10.1111/j.1745-6924.2008.00088.x
- Piccinelli, M., & Wilkinson, G. (2000). Gender differences in depression: Critical review. *British Journal of Psychiatry*, 177, 486–492. doi:10.1192/bjp.177.6.486
- Regier, D. A., Narrow, W. E., Kuhl, E. A., & Kupfer, D. J. (Eds.). (2011). *The conceptual evolution of DSM-5*. Arlington, VA: American Psychiatric Publishing, Inc.
- Shear, K., Halmi, K. A., Widiger, T. A., & Boyce, C. (2007). Sociocultural factors and gender. In W. E. Narrow, M. B. First, P. J. Sirovatka, & D. A. Regier (Eds.), *Age and gender considerations in psychiatric diagnosis: A research agenda for DSM-V* (pp. 65–79). Washington, DC: American Psychiatric Association.
- Slade, T., & Watson, D. (2006). The structure of common DSM-IV and ICD-10 mental disorders in the Australian general population. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*, 36, 1593–1600. doi:10.1017/S0033291706008452
- Slutske, W. S., Heath, A. C., Madden, P. A., Bucholz, K. K., Statham, D. J., & Martin, N. G. (2002). Personality and the genetic risk for alcohol dependence. *Journal of Abnormal Psychology*, 111, 124–133. doi:10.1037/0021-843X.111.1.124
- Trull, T. J., Jahng, S., Tomko, R. L., Wood, P. K., & Sher, K. J. (2010). Revised NESARC personality disorder diagnoses: Gender, prevalence, and comorbidity with substance dependence disorders. *Journal of Personality Disorders*, 24, 412–426. doi:10.1521/pedi.2010.24.4.412
- Vesga-López, O., Schneier, F. R., Wang, S., Heimberg, R. G., Liu, S. M., Hasin, D. S., & Blanco, C. (2008). Gender differences in generalized anxiety disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Journal of Clinical Psychiatry*, 69, 1606–1616.
- Vollebergh, W. A. M., Iedema, J., Bijl, R. V., de Graaf, R., Smit, F., & Ormel, J. (2001). The structure and stability of common mental disorders: The NEMESIS study. *Archives of General Psychiatry*, 58, 597–603. doi:10.1001/archpsyc.58.6.597
- Widiger, T. A. (2007). DSM's approach to gender: History and controversies. In W. E. Narrow, M. B. First, P. J. Sirovatka, & D. A. Regier (Eds.), *Age and gender considerations in psychiatric diagnosis: A research agenda for DSM-V* (pp. 31–46). Washington, DC: American Psychiatric Association.
- Wittchen, H.-U. (1994). Reliability and validity studies of the WHO-Composite International Diagnostic Interview (CIDI): A critical review. *Journal of Psychiatric Research*, 28, 57–84. doi:10.1016/0022-3956(94)90036-1

Received December 22, 2010

Revision received April 27, 2011

Accepted May 3, 2011 ■